



UWS Academic Portal

Benefits of enrichment on the behaviour of ornamental fishes during commercial transport

Vanderzwalmen, Myriam; Carey, Peter; Snellgrove, Donna; Sloman, Katherine A.

Published in:
Aquaculture

DOI:
[10.1016/j.aquaculture.2020.735360](https://doi.org/10.1016/j.aquaculture.2020.735360)

Published: 15/09/2020

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Vanderzwalmen, M., Carey, P., Snellgrove, D., & Sloman, K. A. (2020). Benefits of enrichment on the behaviour of ornamental fishes during commercial transport. *Aquaculture*, 526, [735360].
<https://doi.org/10.1016/j.aquaculture.2020.735360>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Benefits of enrichment on the behaviour of ornamental fishes during commercial transport.

Myriam Vanderzwalmen^a, Peter Carey^b, Donna Snellgrove^c and Katherine A. Sloman^a

^aSchool of Health and Life Sciences, University of the West of Scotland, Paisley, PA1 2BE,
UK

^bPets at Home, Handforth, Cheshire, UK

^cWALTHAM Petcare Science Institute, Waltham-on-the-Wolds, LE14 4RT, Leicestershire,
UK

correspondence

Myriam Vanderzwalmen, School of Health and Life Sciences, University of the West of
Scotland, Paisley PA1 2BE, UK. Tel: 0044 141 8483371 Email:
myriam.vanderzwalmen@uws.ac.uk

Highlights

- The effect of enrichment on fish behaviour during commercial transport was studied.
- Enrichment during transport reduced erratic swimming on arrival and post-transport.
- Enrichment during transport reduced chasing behaviours post-transport.
- Enrichment during transport reduced stress-related behaviours during recovery.
- Enrichment could reduce transport-stress for millions of ornamental fishes.

Abstract

Environmental enrichment is known to reduce stereotypical behaviours in a wide range of captive animals, providing clear welfare benefits in many species. However, one of the most significant stressors faced by captive animals is live transport but whether enrichment can alleviate transport stress is unknown. Using behavioural measures as indicators of fish welfare, we investigated whether the addition of environmental enrichment can improve the welfare of ornamental fish during commercial transport. Pairs of bags containing variatus platy (*Xiphophorus variatus*) were transported by road with or without enrichment (plastic loops) from a UK wholesaler to one of two stores, the first and the last store on a delivery route. Transport time to the first store was short (< 4 h) and longer (> 6 h) to the second. Fish behaviour and incidences of mortality, disease and injury were recorded immediately following transport and during a four-week recovery period. Immediately post-transport, significantly fewer occurrences of erratic swimming were observed in the enriched group compared to the control group, and lower levels of chasing were recorded in the enriched group during recovery. This study is the first to demonstrate the behavioural benefits of enrichment during live transport of fishes under commercial conditions.

Keywords: *environmental enrichment, ornamental fishes, stress, transport, welfare.*

1. Introduction

Enrichment is often added to the housing of captive animals with the aim of improving their welfare (Carlstead and Shepherdson, 2000; Young, 2003). Enrichment is defined as an increase in environmental complexity that aims to reduce maladaptive and aberrant traits in animals (Näslund and Johnsson, 2016). Structural complexity can be added to provide enclosures mimicking the natural environment (Young, 2003), social complexity can enrich an animal's environment through housing with conspecifics or different species (Bloomsith and Lambeth, 2000; De Rouck et al. 2005; Sloman et al. 2011) and dietary complexity can include food flavour, texture and shape variation (Skibieli et al. 2007; Rozek and Millam, 2011). Reported benefits of enrichment include improved feeding, lowered metabolic rates, positive behavioural changes and improved cognitive performance (Millidine et al. 2006; Rozek and Millam, 2011; Mileva and Bielajew, 2015; Grimberg-Henrici et al. 2016). While diversity in the environmental requirements of animals means that the response to enrichment is often species (Costa et al. 2018), life-stage (Ullah et al. 2017) and context-specific (Wood, 1998), positive effects of enrichment have been documented across a wide range of animals (Wood, 1998; Belz et al. 2003; Baumans et al. 2005; Milgram et al. 2005; De Rouck et al. 2005; DeVries et al. 2007; Ellis, 2009; Cornale et al. 2015; Sullivan et al. 2016)

For many captive animals, one of the major challenges to their welfare is the stress they encounter during live transport (Grandin, 1997; Harmon, 2009; Nomura et al. 2009). Given the benefits of enriched housing, it is possible that enriched transport could provide similar effects. Studies on the benefits of enrichment during transport are restricted to pigs and monkeys (Peeters and Geers, 2006; Fernström et al. 2008; Roldan-Santiago et al. 2015);

while these studies have found mixed results there is some evidence to suggest benefits of enrichment during transport. For example, providing maize-filled balls to pigs during transport and lairage resulted in lower plasma lactate concentrations and reduced incidents of shoulder injuries compared to control pigs (N.B. significance was taken as $p < 0.1$) (Peeters and Geers, 2006). Addition of straw bedding during piglet transport attenuated physiological effects of transport stress (Roldan-Santiago et al. 2015). Social enrichment during simulated transport of cynomolgus monkeys (*Macaca fascicularis*) reduced stress-associated behaviours (Fernström et al. 2008).

Ornamental fishes within the aquarium trade experience long transports (from a few hours to >48 h) by road and aeroplanes. Stressors associated with transport include handling, crowding and degrading water quality. The aim of the present study was to assess potential benefits of adding structural enrichment during commercial transport of ornamental fishes. We hypothesised that structural enrichment would provide shelter and disrupt physical contact between fish during transport, leading to reduced stress-related behaviours post-transport. The behaviour of variatus platy (*Xiphophorus variatus*), transported by road with or without enrichment (loops of floating plastic), was examined immediately post-transport and during four weeks of recovery at a retailer.

2. Methods

2.1 Study animals and data collection

Variatus platy (random mix of males and females) were transported from a UK wholesaler to two retail stores (journey range: 8.5-10.7 km). Transportation time to the first store was ~2 h 30 min (short transport) and 7 h to the second store (long transport). At the

wholesaler, platys (~3.5 cm in length) were netted and placed into polythene bags containing tank water (5-10 fish in 1 l, 25.5 x 35 cm bags, bags were ~25% filled). The air at the top of the bags was replaced with pure oxygen and the bags air-tight sealed and placed into an insulated packing box along with a variable number of other bags. The number of fishes in the bags varied based on the orders that were placed by the stores. Twelve bags containing environmental enrichment (five plastic loops per bag, 31.3 cm in diameter and 1.8 ± 0.32 cm in width; mean \pm SD, Fig. 1) and 12 control bags without enrichment were tracked, with six bags of each treatment going to each store. On each journey, bags were tracked in pairs (one control, one enriched), and each pair was assigned a number for statistical analyses accounting for different journey times dependent on traffic delays. Pairs of bags tracked together always contained the same number and colour strain of fish. However, different pairs of bags contained different colour strains to ensure that the fish from different shipments could be identified. At store 1 (short transport) all the fish from one bag were sold prior to the end of the data collection, so this pair was excluded from statistical analysis.




Figure 1: Enrichment. Example photograph of plastic loop used as enrichment during transport (indicated by the arrow).

On arrival at the retail stores, bags were floated in separate recovery tanks for 30 min, during which time fish were recorded using a GoPro hero 5 camera set to 1080P (progressive scan) 120 fps (frames per seconds) with a wide angle. Cameras were mounted on a monopod attached to the outside of the tank. The bags were then opened to allow for slow mixing of the water and after a further 30 min the fish were released into the tank. Staff at the retail stores were unaware of the treatments or study aims to avoid any bias in daily husbandry practices. Preliminary trials found that water parameters in the bags on arrival at the stores were pH: 6.3 ± 0.34 ; oxygen: $188.65\% \pm 23.41$; and temperature: $23.5^{\circ}\text{C} \pm 1.22$ (mean \pm SD), and were not significantly different between transport times ($P > 0.1$ all parameters). Ammonia and nitrate remained at safe levels (i.e. total ammonia nitrogen < 0.93 ppm, unionised ammonia < 0.004 ppm, nitrate < 21.2 ppm) during transport and recovery.

During recovery, fish behaviour was recorded once every 7 days for 4 weeks at random times of the day, but not within 1 h of feeding. The duration of recordings and the camera set up were the same as on arrival. Videos were subsequently used to analyse fish behaviour (Table 1) using Behavioural Observation Research Interactive Software (BORIS) (Friard and Gamba, 2016). Randomly selected focal fish were tracked for 3 min, repeated consecutively from the start to the end of the video such that 10 individuals were tracked. Behaviours for each focal fish were then added to obtain a bag score. Due to the nature of the enrichment analysing the videos blind was not possible as the enrichment was visible in the videos recorded on arrival. Therefore, a subsample (~10%) of videos were analysed blind by an additional observer (inter-observer reliability: Kendall's W coefficient: 0.99). From the videos, any visible body and fin injuries were recorded for focal fish and given a severity score from 0 – 3 where 0 = no injuries and 3 = major injury (Vanderzwalmen et al. 2019). Scores for fin and body injuries were summed to obtain a total injury score recorded for each

123 fish which were then summed to obtain a total injury score per bag. This score was then
124 adjusted for the number of fish that remained at each data collection point to account for fish
125 that had been sold or died. From arrival of the fish until the end of the 4-week recovery
126 period, mortality and occurrence of disease was recorded daily by the retail store staff. All
127 fish were fed a commercial diet *ad libitum* twice daily and kept on a 12 h light/dark regime.
128 Preliminary trials found that water parameters in the tanks at the stores were pH: 6.96 ± 0.51 ;
129 oxygen: $101\% \pm 1.93$; temperature: $22.42^{\circ}\text{C} \pm 1.6$ (mean \pm SD), and were not significantly
130 different between stores ($p > 0.1$ all parameters).

131 Table 1. Ethogram of the behaviours recorded.

Behaviour	Description	Relation to welfare
Biting	Fish biting or nipping a conspecific.	Biting can cause injuries and may lead to mortality (Pitcher, 1986; Noble et al. 2012).
Chases	Occurrence of a fish chasing a conspecific.	Aggressive interaction has been associated with elevated stress (Vera Cruz and Brown, 2007; Kadry and Barreto, 2010; Gronquist and Berges, 2013).
Erratic swimming	Occurrence of rapid swimming and direction change in the absence of being chased.	Erratic swimming is an indicator of elevated stress, distress or pathogenic condition and may be used as a sign of poor welfare (Conte, 2004).
Ventilation rate	Frequency per minute (Alvarenga and Volpato, 1995).	Ventilation rate can increase as a result of stressors and is a highly sensitive mechanism involved in stress coping strategies (Barreto and Volpato, 2004; Martins et al. 2012).

2.2 Statistical analysis

R 3.6.1 was used for statistical analyses (Bolker et al. 2009; R Core Team, 2019). One outlier from the bite data was removed as it was 22.5 SD away from the mean (Miller, 1991). General linear mixed models (GLMM) were carried out on ventilation rate using the lme function from the lme4 package (Bates et al. 2015). Treatment, recovery stage and transport duration were set as explanatory variables; strain had no significant effect on behaviour and was included as a random variable, as was pair number. Number of fish per bag was five in all cases except in one pair where 10 fish per bag were ordered that week. Therefore, number of fish was not included as a random variable due to the low variation causing singularity when included in models. The interactions between the explanatory variables were also analysed. The distribution of residuals was determined visually (R Core Team, 2019). For response variables recorded as counts, a generalised linear mixed model (GLMM) using poisson family from the lme4 package was used with the same variables as above (Bates et al. 2015). As necessary, data were log or square root transformed to obtain normally distributed residuals. Percentage data were expressed as proportions and a binomial GLMM was carried out with the fixed and random variables as described above. The best fit model was selected by disregarding explanatory variables that did not significantly improve the fit of the model (Bates et al. 2015). The significance of the explanatory variables was tested using ANOVA at $P < 0.05$ (Fox and Weisberg, 2011; Pinheiro et al. 2018) and pair-wise post-hoc Tukey tests were carried out (Lenth, 2017). Figures were created using ggplot2 (Wickham et al. 2016).

2.3. Ethics

This study was approved by the University of the West of Scotland Ethics Committee as well as the Animal Welfare Ethics Review Board at the Waltham Petcare Science Institute.

3. Results

Erratic swimming was significantly affected by treatment, recovery stage and transport duration (Table 2). Erratic swimming was consistently lower in fish transported with enrichment than controls (Fig. 2). In fish that experienced the short transport, erratic swimming decreased during week 1 of recovery but then rose within the following weeks to similar levels seen on arrival (Fig. 2). In fish from the long transport, however, erratic swimming was lower during recovery than on arrival and remained low throughout recovery (Fig. 2).

Table 2 ANOVA result of best fit model for erratic swimming and chasing with *F* values, degrees of freedom (df), and *P* values. Significant p-values are indicated in bold.

Behaviour	Fixed Effect	numDF	denDF	<i>F</i> -value	<i>P</i>
Erratic swimming	Treatment	1	78	125.8	<0.0001
	Recovery Stage	4	78	108.31	<0.0001
	Transport Duration	1	9	36.92	<0.0001
	Treatment * Recovery Stage	4	78	2.413	0.075
	Recovery Stage * Transport Duration	4	78	48.062	<0.0001
	Treatment * Transport Duration	1	78	0.391	0.563
	Treatment * Recovery Stage * Transport Duration	4	78	2.404	0.046
Chasing	Treatment	1	88	173.614	<0.0001
	Recovery Stage	4	88	35.09	<0.0001
	Transport Duration	1	5	0.008	0.842
	Treatment * Recovery Stage	4	88	16.184	<0.0001
	Treatment * Transport Duration	1	95	0.196	0.669




Figure 2: Occurrence of erratic swimming in fish transported with or without environmental enrichment for a short and long transport. Data are mean (diamond), median, upper and lower 25th percentile, and outliers, n=5 bags per treatment for short and n=6 for long transport. Asterisks indicate a significant difference between treatments within a time point (post-hoc Tukey, $P < 0.05$). Lowercase letters indicate significance between stage of transport within control (a-f) and enriched (v-z) treatments, with separate post-hoc analyses used for short and long transport duration. Boxes sharing a letter are not significantly different (post-hoc Tukey, $P < 0.05$).

Chasing was significantly affected by treatment and recovery stage with a significant interaction between the two factors (Table 2). However, unlike for erratic swimming, there was no effect of transport duration on chasing. Chasing behaviour was low on arrival at the stores for both treatments but increased during the recovery period, remaining lower in the fish transported with enrichment (Fig. 3). In fish from the control treatment, chasing was significantly higher than on arrival by week 1 of recovery and remained high, while the number of chases in the enriched treatment only began to increase slowly from week 3 of recovery (Fig. 3). Biting behaviour, injury score, mortality and ventilation rate were not significantly affected by treatment, recovery stage or duration ($P > 0.05$ for all, Table S1 supplementary material) and remained low throughout (Table S2 supplementary material). Occurrence of disease was too low to carry out statistical analyses.




Figure 3: Occurrence of chasing in fish transported with or without environmental enrichment. Data are mean (diamond), median, upper and lower 25th percentile, and outliers, n=11. Asterisks indicate a significant difference between treatments within a specific time point (post-hoc Tukey, $P < 0.05$). Letters indicate a significant difference between stages of transport within control (a-c) and enriched (x,y) treatments where boxes sharing a letter are not significantly different (post-hoc Tukey, $P < 0.05$).

4. Discussion

Fish that were transported with enrichment showed a reduction in stress-associated behaviours including erratic swimming and chasing. Erratic swimming in the enriched group was lower on arrival at stores than in the control group, and during recovery fish shipped with enrichment showed lower levels of chasing.

Some chasing and biting are expected during hierarchy establishment, but prolonged chasing can lead to chronic stress for the fish subjected to the aggressive behaviour (Braddock, 1945; Scott and Currie, 1980). Following stress, fish can display displaced aggression behaviours, such as chasing, as a coping mechanism to manage their own stress (Sneddon et al., 2016). Reduced aggression could therefore indicate improved stress-coping capabilities in the enriched group (Sneddon et al., 2016). Lower levels of chasing in the enriched group may also suggest a less aggressive establishment of hierarchy. Biting behaviours are a greater welfare concern than chasing due to risks of injury and infection but were low throughout the present study.

Erratic swimming was higher following the short transport compared to the long transport, potentially as the longer time that elapsed between packing and unpacking allowed greater recovery from handling on route for the long transport fish. This hypothesis is supported by previous studies on fish transport which have found that handling is the most stressful part of transport and recovery begins within the first 2-4 h of transport duration (Pickering et al. 1982; Acerete et al. 2004; Nomura et al. 2009). The majority of studies that have considered the welfare of ornamental fishes during transport have used simulated transport and focussed on stress recovery for 24-96 h post-transport (Zanuzzo et al. 2012; Ramírez-duarte et al. 2013; Salaro et al. 2015; Pattanasiri et al. 2017). However, as animals

can experience stress for extended periods of time following transport (Knowles et al. 1993; Dembiec et al. 2004; Fernström et al. 2008), in the present study indicators of stress were recorded for four weeks post-transport. While differences in behaviour between treatments were seen over this time frame, mortality, disease and injury were not affected by enrichment most likely as levels of all three remained low throughout the study.

Although a wealth of literature exists on the use of environmental enrichment for fish holding conditions (Näslund and Johnsson, 2016), to the authors' knowledge, no previous studies have looked at the potential benefits of enrichment for fishes during commercial transport. However, there are a few studies that have considered benefits of enrichment during periods of acute stress. For example, zebrafish (*Danio rerio*) from enriched tanks released less cortisol than zebrafish from bare tanks following a pursuit stress (Giacomini et al. 2016) and a similar result was seen in jundiá (*Rhamdia quelen*) (Barcellos et al. 2009). Exposure to blue substrate prior, during and after a confinement stress also improved the recovery rate of serotonin and plasma glucose in gilthead seabream (*Sparus aurata*) (Batzina et al. 2014). The effects of enrichment during transport have been considered in other animals showing positive effects of social enrichment in cynomolgus monkeys (Fernström et al. 2008) and the addition of toys (Peeters and Geers, 2006) and substrate (Roldan-Santiago et al. 2015) for pigs.

Plastic loops were chosen as the enrichment as they are already used within the ornamental trade to reduce the stress experienced by some species, meaning it is an enrichment which reflects current practice and could be implemented for other species. Prior to the present study, there had been no research on the use of plastic loops as enrichment

during ornamental fish transport. Plastic loops are likely to disturb visual contact between fishes within the transport bag and might provide hiding places, however, further research is needed to better understand the functionality of the plastic loops as enrichment. More research is also needed to identify other potential forms of enrichment during ornamental fish transport across different species and life-stages. Consideration must be given to the fact that the type of environmental enrichment provided is likely to affect both the response of the animal and any beneficial effects which has been shown in non-transport related studies. For example, goldfish (*Carassius auratus*) displayed increased vigilance in the presence of a predator when a solid block was used as enrichment but not when artificial plants were available (Ingrum et al. 2010). Black bucks (*Antelope cervicapra*) were found to prefer a tree trunk/branch combination as enrichment compared to vertical brushes (Bono et al. 2016) and cattle (Charolais cross heifers) prefer scratching enrichments over olfactory enrichments (Wilson et al. 2002). Goldfish also prefer enriched areas over non-enriched areas with no preference between artificial or real plants indicating that the functionality of enrichment can be more important than the material it is made from (Sullivan et al. 2016).

5. Conclusion

Millions of animals are transported daily across the world. Transport is stressful for these animals, yet surprisingly few studies have assessed the effect of enrichment during animal transport. This study is the first to use enrichment during the commercial transport of ornamental fishes, finding a clear benefit in the reduction of stress-associated behaviours. With growing pressure on industries to improve the welfare of the animals they care for, further research into environmental enrichment to manage stress during commercial transport is warranted.

Acknowledgments

Many thanks to management and staff at the wholesaler and retail stores where data were collected.

6. References

- Acerete, L., Balasch, J.C., Espinosa, E., Josa, A., Tort, L., 2004. Physiological responses in Eurasian perch (*Perca fluviatilis*, L.) subjected to stress by transport and handling. *Aquaculture* 237, 167–178. <https://doi.org/10.1016/j.aquaculture.2004.03.018>
- Alvarenga, C.M.D., Volpato, G.L., 1995. Agonistic profile and metabolism in alevins of the Nile Tilapia. *Physiol. Behav.* 57, 75–80. [https://doi.org/10.1016/0031-9384\(94\)00206-K](https://doi.org/10.1016/0031-9384(94)00206-K)
- Barcellos, L.J.G., Kreutz, L.C., Quevedo, R.M., da Rosa, J.G.S., Koakoski, G., Centenaro, L., Pottker, E., 2009. Influence of color background and shelter availability on jundiá (*Rhamdia quelen*) stress response. *Aquaculture* 288, 51–56. <https://doi.org/10.1016/j.aquaculture.2008.11.002>
- Barreto, R.E., Volpato, G.L., 2004. Caution for using ventilatory frequency as an indicator of stress in fish. *Behav. Processes* 66, 43–51. <https://doi.org/10.1016/j.beproc.2004.01.001>
- Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Batzina, A., Kalogiannis, D., Dalla, C., Papadopoulou-Daifoti, Z., Chadio, S., Karakatsouli, N., 2014. Blue substrate modifies the time course of stress response in gilthead seabream *Sparus aurata*. *Aquaculture* 420–421, 247–253. <https://doi.org/10.1016/j.aquaculture.2013.11.016>

- 291 Baumans, V., 2005. Environmental enrichment for laboratory rodents and rabbits:
292 Requirements of rodents, rabbits, and research. *ILAR J.* 46, 162–170.
293 <https://doi.org/10.1093/ilar.46.2.162>
- 294 Belz, E.E., Kennell, J.S., Czambel, R.K., Rubin, R.T., Rhodes, M.E., 2003. Environmental
295 enrichment lowers stress-responsive hormones in singly housed male and female rats.
296 *Pharmacol. Biochem. Behav.* 76, 481–486. <https://doi.org/10.1016/j.pbb.2003.09.005>
- 297 Bloomsmith, M.A., Lambeth, S.P., 2000. Videotapes as enrichment for captive chimpanzees
298 (*Pan troglodytes*). *Zoo Biol.* 19, 541–551.
- 299 Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H.,
300 White, J.S.S., 2009. Generalized linear mixed models: a practical guide for ecology and
301 evolution. *Trends Ecol. Evol.* 24, 127–135. <https://doi.org/10.1016/j.tree.2008.10.008>
- 302 Bono, L., Mongillo, P., De Boni-Russo, G., Gabai, G., Normando, S., 2016. Effects of two
303 forms of environmental enrichment on a group of captive blackbucks (*Antilope cervicapra*):
304 A pilot study. *J. Vet. Behav. Clin. Appl. Res.* 12, 66–72.
305 <https://doi.org/10.1016/j.jveb.2016.02.003>
- 306 Braddock, J.C., 1945. Some aspects of the dominance-subordination relationship in the fish
307 *Platypleurodon maculatus*. *Physiol. Zool.* 18, 176–195.
308 <https://doi.org/10.1086/physzool.18.2.30151863>
- 309 Carlstead, K., Shepherdson, D., 2000. Alleviating stress in zoo animals with environmental
310 enrichment, in: Moberg, G.P., Mench, J. (Eds.), *The Biology of Animal Stress: Basic*
311 *Principles and Implications for Animal Welfare*. CABI publishing, Oxford, UK, pp. 337–354.
- 312 Conte, F., 2004. Stress and the welfare of cultured fish. *Appl. Anim. Behav. Sci.* 86, 205–
313 223. <https://doi.org/10.1016/j.applanim.2004.02.003>

- 314 Cornale, P., Macchi, E., Miretti, S., Renna, M., Lussiana, C., Perona, G., Mimosi, A., 2015.
- 315 Effects of stocking density and environmental enrichment on behavior and fecal
- 316 corticosteroid levels of pigs under commercial farm conditions. *J. Vet. Behav. Clin. Appl.*
- 317 *Res.* 10, 569–576. <https://doi.org/10.1016/j.jveb.2015.05.002>
- 318 Costa, R., Sousa, C., Llorente, M., 2018. Assessment of environmental enrichment for
- 319 different primate species under low budget: A case study. *J. Appl. Anim. Welf. Sci.* 21, 185–
- 320 199. <https://doi.org/10.1080/10888705.2017.1414606>
- 321 De Rouck, M., Kitchener, A.C., Law, G., Nelissen, M., 2005. A comparative study of the
- 322 influence of social housing conditions on the behaviour of captive tigers (*Panthera tigris*).
- 323 *Univ. Fed. Anim. Welf.* 14, 229–238.
- 324 Dembiec, D.P., Snider, R.J., Zanella, A.J., 2004. The effects of transport stress on tiger
- 325 physiology and behavior. *Zoo Biol.* 23, 335–346. <https://doi.org/10.1002/zoo.20012>
- 326 DeVries, T.J., Vankova, M., Veira, D.M., von Keyserlingk, M.A.G., 2007. Short
- 327 communication: Usage of mechanical brushes by lactating dairy cows. *J. Dairy Sci.* 90,
- 328 2241–2245. <https://doi.org/10.3168/jds.2006-648>
- 329 Ellis, S., 2009. Environmental enrichment: practical strategies for improving feline welfare.
- 330 *J. Feline Med. Surg.* 11, 901–912.
- 331 Fernström, A.L., Sutian, W., Royo, F., Westlund, K., Nilsson, T., Carlsson, H.E., Paramastri,
- 332 Y., Pamungkas, J., Sajuthi, D., Schapiro, S.J., Hau, J., 2008. Stress in cynomolgus monkeys
- 333 (*Macaca fascicularis*) subjected to long-distance transport and simulated transport housing
- 334 conditions. *Stress* 11, 467–476. <https://doi.org/10.1080/10253890801903359>
- 335 Fox, J., Weisberg, S., 2011. *An R Companion to Applied Regression*, second. ed. Sage,
- 336 Thousand Oaks, CA.

- 337 Friard, O., Gamba, M., 2016. BORIS: a free, versatile open-source event-logging software for
338 video/audio coding and live observations. *Methods Ecol. Evol.* 7, 1325–1330.
339 <https://doi.org/10.1111/2041-210X.12584>
- 340 Giacomini, A.C.V.V., Abreu, M.S., Zanandrea, R., Saibt, N., Friedrich, M.T., Koakoski, G.,
341 Gusso, D., Piato, A.L., Barcellos, L.J.G., 2016. Environmental and pharmacological
342 manipulations blunt the stress response of zebrafish in a similar manner. *Sci. Rep.* 6, 1–6.
343 <https://doi.org/10.1038/srep28986>
- 344 Grandin, T., 1997. Assessment of stress during handling and transport. *J. Anim. Sci.* 75, 249–
345 257.
- 346 Grimberg-Henrici, C.G.E., Vermaak, P., Bolhuis, E.J., Nordquist, R.E., van der Staay, F.J.,
347 2016. Effects of environmental enrichment on cognitive performance of pigs in a spatial
348 holeboard discrimination task. *Anim. Cogn.* 19, 271–283. [https://doi.org/10.1007/s10071-](https://doi.org/10.1007/s10071-015-0932-7)
349 [015-0932-7](https://doi.org/10.1007/s10071-015-0932-7)
- 350 Gronquist, D., Berges, J.A., 2013. Effects of aquarium-related stressors on the zebrafish: A
351 comparison of behavioral, physiological, and biochemical indicators. *J. Aquat. Anim. Health*
352 25, 53–65. <https://doi.org/10.1080/08997659.2012.747450>
- 353 Harmon, T.S., 2009. Methods for reducing stressors and maintaining water quality associated
354 with live fish transport in tanks: a review of the basics. *Rev. Aquac.* 1, 58–66.
355 <https://doi.org/10.1111/j.1753-5131.2008.01003.x>
- 356 Ingrum, J., Nordell, S.E., Dole, J., 2010. Effects of habitat complexity and group size on
357 perceived predation risk in goldfish (*Carassius auratus*). *Ethol. Ecol. Evol.* 22, 119–132.
358 <https://doi.org/10.1080/03949371003707638>
- 359 Kadry, V.O., Barreto, R.E., 2010. Environmental enrichment reduces aggression of pearl

- 360 cichlid, *Geophagus brasiliensis*, during resident-intruder interactions Vivian. *Neotrop.*
361 *Ichthyol.* 8, 329–332. <https://doi.org/10.1590/S1679-62252010000200011>
- 362 Knowles, T.G., Warriss, P.D., Brown, S.N., Kestin, S.C., Rhind, S.M., Edwards, J.E., Anil,
363 M.H., Dolan, S.K., 1993. Long distance transport of lambs and the time needed for
364 subsequent recovery. *Vet. Rec.* 133, 283–286.
- 365 Lenth, R. V, 2017. Least-squares means: The R package lsmeans. *J. Stat. Softw.* 69, 1–33.
366 <https://doi.org/10.18637/jss.v069.i01>
- 367 Martins, C.I.M., Galhardo, L., Noble, C., Damsgård, B., Spedicato, M.T., Zupa, W.,
368 Beauchaud, M., Kulczykowska, E., Massabuau, J.-C., Carter, T., Planellas, S.R., Kristiansen,
369 T., 2012. Behavioural indicators of welfare in farmed fish. *Fish Physiol. Biochem.* 38, 17–41.
370 <https://doi.org/10.1007/s10695-011-9518-8>
- 371 Mileva, G.R., Bielajew, C., 2015. Environmental manipulation affects depressive-like
372 behaviours in female Wistar-Kyoto rats. *Behav. Brain Res.* 293, 208–216.
373 <https://doi.org/10.1016/j.bbr.2015.07.035>
- 374 Milgram, N.W., Head, E., Zicker, S.C., Ikeda-Douglas, C.J., Murphey, H., Muggenburg, B.,
375 Siwak, C., Tapp, D., Cotman, C.W., 2005. Learning ability in aged beagle dogs is preserved
376 by behavioral enrichment and dietary fortification: A two-year longitudinal study. *Neurobiol.*
377 *Aging* 26, 77–90. <https://doi.org/10.1016/j.neurobiolaging.2004.02.014>
- 378 Miller, J., 1991. Short report: reaction time analysis with outlier exclusion: bias varies with
379 sample size. *Q. J. Exp. Psychol.* 43, 907–912. <https://10.1080/14640749108400962>
- 380 Millidine, K.J., Armstrong, J.D., Metcalfe, N.B., 2006. Presence of shelter reduces
381 maintenance metabolism of juvenile salmon. *Funct. Ecol.* 20, 839–845.
382 <https://doi.org/10.1111/j.1365-2435.2006.01166.x>

- 383 Näslund, J., Johnsson, J.I., 2016. Environmental enrichment for fish in captive environments:
384 Effects of physical structures and substrates. *Fish Fish.* 17, 1–30.
385 <https://doi.org/10.1111/faf.12088>
- 386 Noble, C., Jones, H.A.C., Damsgård, B., Flood, M.J., Midling, K.O., Roque, A., Sæther, B.S.,
387 Cottee, S.Y., 2012. Injuries and deformities in fish: Their potential impacts upon aquacultural
388 production and welfare. *Fish Physiol. Biochem.* 38, 61–83. [https://doi.org/10.1007/s10695-](https://doi.org/10.1007/s10695-011-9557-1)
389 [011-9557-1](https://doi.org/10.1007/s10695-011-9557-1)
- 390 Nomura, M., Sloman, K.A., von Keyserlingk, M.A.G., Farrell, A.P., 2009. Physiology and
391 behaviour of Atlantic salmon (*Salmo salar*) smolts during commercial land and sea transport.
392 *Physiol. Behav.* 96, 233–243. <https://doi.org/10.1016/j.physbeh.2008.10.006>
- 393 Pattanasiri, T., Taparhudee, W., Suppakul, P., 2017. Anaesthetic efficacy of clove oil-coated
394 LDPE bag on improving water quality and survival in the Siamese fighting fish, *Betta*
395 *splendens*, during transportation. *Aquac. Int.* 25, 197–209. [https://doi.org/10.1007/s10499-](https://doi.org/10.1007/s10499-016-0020-2)
396 [016-0020-2](https://doi.org/10.1007/s10499-016-0020-2)
- 397 Peeters, E., Geers, R., 2006. Influence of provision of toys during transport and lairage on
398 stress responses and meat quality of pigs. *Anim. Sci.* 82, 591–595.
399 <https://doi.org/10.1079/asc200686>
- 400 Pickering, A., Pottinger, T., Christie, P., 1982. Recovery of the brown trout, *Salmo trutta* L.,
401 from acute handling stress: a time course study. *J. Fish Biol.* 20, 229–244.
- 402 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2018. nlme: Linear and
403 Nonlinear Mixed Effects Models. <https://cran.r-project.org/web/packages/nlme/index.html>
- 404 Pitcher, T.J. (Ed.), 1986. The Behaviour of Teleost Fishes, Springer Science & Business
405 Media. Croom Helm, London, UK. https://doi.org/10.1007/978-1-4684-8261-4_5

- 406 R Core Team, 2019. R: A Language and Environment for Statistical Computing.
- 407 Ramírez-duarte, W.F., Pineda-quiroya, C., Rueda, N.M., 2013. Evaluation of the use of
 408 sodium chloride, eugenol, and zeolite in confinement of *Ancistrus triradiatus*. *Orinoquia* 17,
 409 84–95.
- 410 Roldan-Santiago, P., Trujillo-Ortega, M., Borderas-Tordesillas, F., Martínez-Rodríguez, R.,
 411 Mora-Medina, P., Flores-Peinado, S., Sánchez-Hernández, M., García-Herrera, R., González-
 412 Lozano, M., Mota-Rojas, D., 2015. Physiometabolic responses to road transport in weaned
 413 piglets for a short period and the effects of straw bedding. *Anim. Sci. J.* 86, 563–571.
 414 <https://doi.org/10.1111/asj.12324>
- 415 Rozek, J.C., Millam, J.R., 2011. Preference and motivation for different diet forms and their
 416 effect on motivation for a foraging enrichment in captive orange-winged Amazon parrots
 417 (*Amazona amazonica*). *Appl. Anim. Behav. Sci.* 129, 153–161.
 418 <https://doi.org/10.1016/j.applanim.2010.11.009>
- 419 Salaro, A.L., Camplelo, D.A.V., Tavares, M.M., Braga, L.G.T., Pontes, M.D., Zuanon,
 420 J.A.S., 2015. Transport of *Astyanax altiparanae* Garutti and Britski, 2000 in saline water.
 421 *Acta Sci. - Biol. Sci.* 37, 137–142. <https://doi.org/10.4025/actascibiolsci.v37i2.26884>
- 422 Scott D.B.C., Currie C.E. 1980 Social hierarchy in relation to adrenocortical activity in
 423 *Xiphophorus helleri* Heckel. *J. Fish Biol.* 16, 265–277. [https://doi.org/10.1111/j.1095-
 424 8649.1980.tb03704.x](https://doi.org/10.1111/j.1095-8649.1980.tb03704.x)
- 425 Skibieli, A.L., Trevino, H.S., Naugher, K., 2007. Comparison of several types of enrichment
 426 for captive felids. *Zoo Biol.* 26, 371–381. <https://doi.org/10.1002/zoo>
- 427 Sloman, K.A., Baldwin, L., McMahon, S., Snellgrove, D., 2011. The effects of mixed-species
 428 assemblage on the behaviour and welfare of fish held in home aquaria. *Appl. Anim. Behav.*

- 429 *Sci.* 135, 160–168. <https://doi.org/10.1016/j.applanim.2011.08.008>
- 430 Sneddon, L.U., Wolfenden, D.C.C., Thomson, J.S., 2016. Stress management and welfare, in:
- 431 Schreck, C.B., Tort, L., Farrell, A.P., Brauner, C.J. (Eds.), *Biology of Stress in Fish: Fish*
- 432 *Physiology*. Elsevier Inc., London, UK, pp. 463–539. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-802728-8.00012-6)
- 433 [802728-8.00012-6](https://doi.org/10.1016/B978-0-12-802728-8.00012-6)
- 434 Sullivan, M., Lawrence, C., Blache, D., 2016. Why did the fish cross the tank? Objectively
- 435 measuring the value of enrichment for captive fish. *Appl. Anim. Behav. Sci.* 174, 181–188.
- 436 <https://doi.org/10.1016/j.applanim.2015.10.011>
- 437 Ullah, I., Zuberi, A., Khan, K.U., Ahmad, S., Thörnqvist, P.O., Winberg, S., 2017. Effects of
- 438 enrichment on the development of behaviour in an endangered fish mahseer (*Tor putitora*).
- 439 *Appl. Anim. Behav. Sci.* 186, 93–100. <https://doi.org/10.1016/j.applanim.2016.10.016>
- 440 Vanderzwalmen, M., Edmonds, E., Carey, P., Snellgrove, D., Sloman, K. A., 2019. Effect of
- 441 a water conditioner on ornamental fish behaviour during commercial transport. *Aquaculture*
- 442 514, 734486
- 443 Vera Cruz, E.M., Brown, C.L., 2007. The influence of social status on the rate of growth, eye
- 444 color pattern and Insulin-like Growth Factor-I gene expression in Nile tilapia, *Oreochromis*
- 445 *niloticus*. *Horm. Behav.* 51, 611–619. <https://doi.org/10.1016/j.yhbeh.2007.02.010>
- 446 Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New
- 447 York.
- 448 Wilson, S.C., Mitlöhner, F.M., Morrow-Tesch, J., Dailey, J.W., McGlone, J.J., 2002. An
- 449 assessment of several potential enrichment devices for feedlot cattle. *Appl. Anim. Behav. Sci.*
- 450 76, 259–265. [https://doi.org/10.1016/S0168-1591\(02\)00019-9](https://doi.org/10.1016/S0168-1591(02)00019-9)
- 451 Wood, W., 1998. Interactions among environmental enrichment, viewing crowds, and zoo

- 1417
1418
1419 452 chimpanzees (*Pan troglodytes*). *Zoo Biol.* 17, 211–230. [https://doi.org/10.1002/\(SICI\)1098-](https://doi.org/10.1002/(SICI)1098-)
1420
1421 453 [2361\(1998\)17:3<211::AID-ZOO5>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1098-2361(1998)17:3<211::AID-ZOO5>3.0.CO;2-C)
1422
1423
1424 454 Young, R.J., 2003. Environmental Enrichment for Captive Animals, 1st ed. Blackwell
1425
1426 455 Science Ltd, Oxford, UK.
1427
1428
1429 456 Zanuzzo, F.S., Biller-Takahashi, J.D., Urbinati, E.C., 2012. Effect of *Aloe vera* extract on the
1430
1431 457 improvement of the respiratory activity of leukocytes of matrinxã during the transport stress.
1432
1433 458 *Rev. Bras. Zootec.* 41, 2299–2302. <https://doi.org/10.1590/S1516-35982012001000023>
1434
1435
1436 459
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475

End Section Statements

Competing Interests

The authors declare no competing interests.

Funding

The work was part-funded by Waltham Petcare Science Institute (Mars Petcare) and the University of the West of Scotland.

Declaration of interests

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Myriam Vanderzwalmen: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft

Peter Carey: Conceptualization, Resources, Writing - Review & Editing

Donna Snellgrove: Conceptualization, Writing - Review & Editing

Katherine Sloman: Conceptualization, Methodology, Writing - Review & Editing, Supervision

1 Supplementary Material

2 Table S1 ANOVA result of best fit model for biting, injury score, mortality and ventilation rate with *F* values,
3 degrees of freedom (df), and *P* values.

	Fixed Effect	numDF	denDF	<i>F</i> -value	<i>P</i>
Biting	Treatment	1	91	3.634	0.55
	Recovery Stage	4	91	0.366	0.83
	Transport Duration	1	9	0.098	0.75
Injury score	Treatment	1	72	0.044	0.833
	Recovery Stage	4	72	0.271	0.846
	Transport Duration	1	5	2.225	0.195
Mortality	Treatment	1	94	2.434	0.122
	Recovery Stage	4	94	0.764	0.551
	Transport Duration	1	5	0.054	0.825
Ventilation Rate	Treatment	1	92	0.003	0.955
	Recovery Stage	4	92	2.16	0.079
	Transport Duration	1	4	0.215	0.661

5 Table S2: Mean (\pm SD) for non-significant welfare indicators. Actual mortality data are not presented due to
6 commercial sensitivity.

Stage	Treatment	Transport duration	Biting (30 min ⁻¹)	Ventilation rate min ⁻¹	Injury score (average per fish, see text)
Arrival	Control	Short	0 (\pm 0)	105.05 (\pm 9.11)	0 (\pm 0)
Arrival	Enriched	Short	0 (\pm 0)	99.70 (\pm 4.71)	0 (\pm 0)
Arrival	Control	Long	2.6 (\pm 5.81)	106.80 (\pm 10.20)	0.04 (\pm 0.09)
Arrival	Enriched	Long	0 (\pm 0)	116.49 (\pm 28.45)	0.08 (\pm 0.18)
Week 1	Control	Short	0 (\pm 0)	105.74 (\pm 5.71)	0 (\pm 0)
Week 1	Enriched	Short	0.80 (\pm 1.30)	98.66 (\pm 7.79)	0.08 (\pm 0.18)
Week 1	Control	Long	0.4 (\pm 0.89)	102.27 (\pm 12.75)	0.267 (\pm 0.60)
Week 1	Enriched	Long	0 (\pm 0)	106.41 (\pm 13.81)	0.32 (\pm 0.46)
Week 2	Control	Short	0.4 (\pm 0.89)	107.54 (\pm 14.60)	0.08 (\pm 0.18)
Week 2	Enriched	Short	0 (\pm 0)	110.57 (\pm 13.93)	0.2 (\pm 0.45)
Week 2	Control	Long	0.5 (\pm 0.55)	106.04 (\pm 13.62)	0.07(\pm 0.10)
Week 2	Enriched	Long	0.17 (\pm 0.41)	105.49 (\pm 13.23)	0.8 (\pm 0.167)
Week 3	Control	Short	0.2 (\pm 0.45)	111.89 (\pm 8.05)	0.28 (\pm 0.39)
Week 3	Enriched	Short	0 (\pm 0)	106.19 (\pm 7.88)	0.04 (\pm 0.09)
Week 3	Control	Long	0.33 (\pm 0.82)	112.83 (\pm 12.63)	0.12 (\pm 0.31)
Week 3	Enriched	Long	0 (\pm 0)	114.65 (\pm 20.74)	0.32 (\pm 0.49)
Week 4	Control	Short	0.2 (\pm 0.45)	106.73 (\pm 10.03)	0 (\pm 0)
Week 4	Enriched	Short	0 (\pm 0)	103.99 (\pm 10.03)	0.1 (\pm 0.22)
Week 4	Control	Long	0.5 (\pm 0.84)	102.41 (\pm 13.44)	0.5 (\pm 0.78)
Week 4	Enriched	Long	0 (\pm 0)	103.58 (\pm 17.01)	0.19 (\pm 0.30)

7